## **U MONOCEROTIS FROM 1982 TO 1992**

**abstract**: visual photometry of U Mon for a period of ten years are presented. The study is based on visual observations obtained by GEOS and AFOEV observers, these last downloaded via anonymous-ftp from the related database.

Introduction: the class of RV Tauri stars comprises little more than hundreds of pulsating semiregular variable stars with unique characteristics of variability. Their light curve display two unequal minima (deep and shallow), with a fundamental period inclusive between 30 and 150 days, and, superposed, for the RVb (as U Mon), long-term periodicities of the order of 1000 days. Spectral types range from F5 to late K with peculiarities (TiO band and Ca II versus hydrogen lines).

The star: U Monocerotis is a RV Tauri (subclass b) variable star located at (2000):

 $\alpha = 07^{h} 30^{m} 47^{s} \delta = -09^{\circ} 46' 30''$ 

Observable simply with a binoculars, the star show a light curve that ranges between magn. +6.1 and magn. +8.8, with a period around 92 days .GCVS 85 gives the following ephemer's 2438496+91.32E.

The spectrum ranges from F8EVIB at the maximum of light curve and K0PIB at the minimum.

The theoretical Visual Absolute Magnitude M<sub>V</sub> for a RV Tauri is given by the formula:

 $(1) < M_V > = -5.3 + 0.021P^d$ 

the period P is expressed in days.

The behaviour of the star is well know; purpose in fact of this study is to monitor the fundamental and the long-term periodes.

So I've collected visual observations obtained by GEOS members; members that has sent me your data after the "Appel de Mesures" appeared in NC 632 and 645.

To cover any gaps (small number of observations), I've downloaded via anonymous-ftp from the AFOEV Database at CDS the complete file regarding U Monocerotis<sup>1</sup>.

For details see tables 1 (observations per year) and 2 (number of observations per observer).

Fig. 1 shows a complete light curve of U Mon from 1982 to 1992; look at the two fundamental variations: the first seasonal with a period around 90 days, and the second with a

period around 2500 days (the graph show two minima); the amplitude of range is decreasing versus the minimum and increasing after, a RV Tauri subclass b peculiarity well observable in the light curve of U Mon.





Fig. 2 show the result of a Time Serie Analysis, operated with the program HSH developed by Gaspani.




The two peaks signed with A and B at (respectly) v=0.0103 and v=0.0217 give the values for the period ( $v_A$ ) and the semi-period ( $v_B$ ); the first great peak at v=0.0004 gives a value  $\cong 2.500$  days, corresponding to the long-term variation; others are assumed as noise.


For  $v_A$  the resulting period is 97.08 days; for  $v_B$  46.08 days.

Figures 3,4,5,6,7,8,9,10,11,12 show the light curve of U Mon at every season, respectly

<sup>&</sup>lt;sup>1</sup> Ftp.cdsarc.u-strasbg.fr/pub/afoev/u

from 1982/83 to 1991/92; the points are the single observations, the line represents the best fit.





Conclusion: observing figure 2, we can infer that the period of U Mon is increasing.

More observations are necessary to monitor and to determine the period, above all for the long-term variation.

U Mon

Using formula (1),we can obtain the Visual Absolute Magnitude too :  $<M_V>\sim$  -3.30 ,a considerable value!!

Acknowledgments: I would like to thank everyone who sent me their observations, in detail Mr. Mario Checcucci (CHC).

A special thank to Mr. Emile Schweitzer, who sent me the AFOEV ftp address.

## Maurizio Bignotti

	rabie r								:		
season n. obs	82/83 72	83/84 88	84/85 121	85/86 85	86/87 190	87/8 140		88/89 126	89/90 169	90/91 518	91. 3!
	Table 2										
	AFOEV	PVH	CHC	DMT	BGN 137	FRL 68	BAR 30	IELO 23	FDD 9		

## Bibliography

- 1. M. Bignotti-GEOS NC 788
- 2. DuPUY-Astroph. J. 185
- 3. Gehrz-Astroph. J. 178